

**Removal Lead (Pb) And Mercury (Hg) From Juaro Fish (*Pangasius polyuranodon*) Using Citric Acid From Pineapple Extract (*Ananas comosus*) as Chelating Agent**Pra Dian Mariadi<sup>1</sup>, Rosnita Sebayang<sup>1</sup><sup>1</sup>Faculty of Health Science, Musi Charitas Catholic University, Palembang, IndonesiaCorresponding author e-mail : [pradian\\_dian@yahoo.co.id](mailto:pradian_dian@yahoo.co.id)**ABSTRACT**

An assessment of dietary risk of heavy metals exposure to human is important since it is the main of exposure. The aim of study to measure the contamination of lead and mercury in juaro fish meat and the effort to reduce contaminations using citric acid from pineapple extract as a chelating agent. Samples was soaked and boiled in citric acid extract at concentration 50%, 75 %, 100 % for, 15 min, 30 min, 45 min at 25 °C, 50 °C and 100 °C. The concentration of lead and mercury in Juaro fish lower than the maximum acceptable lever for Pb and Hg respectively (1 mg/kg for Pb, 0.5 mg/Kg for Hg). The result indicated that after soaking and boiling in citric acid solution form pineapple extract at concentration 100 % for 45 min at 100 °C reducing heavy metals Pb from 0.02 mg.Kg<sup>-1</sup> to 0.003 mg.Kg<sup>-1</sup> and Hg from 0.011 mg.Kg<sup>-1</sup> to lower than 0.0001 mg.Kg<sup>-1</sup>. The increasing of soaking time and boiling temperature, the levels of Pb and Hg in Juaro fish Tissue will decreasing.

*Key words: Citric Acid solution, Pineapple extract, heavy metal***INTRODUCTION**

Heavy metals are the most important and basic pollutants because of their bioaccumulation and high toxicity even at very low dosage and cause serious dangers to ecological system as well as human health (Tadjuradi, 2016). With the rapid growth of population and industrialization in the world, the frequency of accidently heavy metal pollution in soil, food and aquatic ecosystems boosted fast in recent years. Due to its non-biodegradation and non decomposing characters. Heavy metals ion can be accumulated through food chain and absorbed by human being.

Pollutions in the aquatic ecosystems has received much concern, due to the abundance, toxicity, persistence, ubiquity and non-degradability in the ecosystems and subsequent bio-accumulation of heavy metals (Liu *et al*, 2016). Heavy metals in polluted habitats may accumulate especially in fish tissue which in turn may enter into human food chain and eventually lead health risk (Ahmed *et al*, 2015). Eddy *et al* (2012) had reported the aquatic ecosystems in musi river contaminated by heavy metals such as lead and mercury. Juaro Fish (*Pangasius polyuranodon*) is one of aquatic species which is thought to have high lead and mercury accumulations due to predatory fish.

Mercury is one the most hazardous heavy metal contami-

nants. The environment pollution by mercury originates from two source-natural and anthropogenic. Coal burning, waste combustion, mining, sewage sludge, deposition, Hg wood impregnation and industrial practices are the main anthropogenic sources. (Alloway, 1995). Long term exposure to mercury compounds from food and water can lead to toxic effects on brain, cardiovascular, pulmonary, urinary, gastrointestinal and neurology system (Kim *et al*, 2016; Turaga *et al*, 2014). Lead causes severe health risks such as fatigue, irritability, myalgia, coma, kidney, liver and brain damage, seizures encephalopathy nervous system dysfunction, disturbance of immune system and development of cancer (Sander *et al*, 2009; Goldstein *et al*, 1992).

The technologies used for reduce contaminations in food using chelating agent. Citric acid has potential as chelating agent to remove toxic metals (Jiang *et al*, 2017; Aderhold *et al*, 2017; Shakkour *et al*, 2014). Pineapple (*Ananas comosus*) is one of the fruits that have high concentrations of citric acid. Citric acid includes a chelating agent which is a stabilizer at food processing. Citric acid binds the heavy metals in the form of complex bond and can reduce side effects of heavy metals in food (Hikmawati and Iliis, 2016).

Interaction of citric acid with heavy metals can be enhanced by soaking (Ulfa *et al*, 2014) and boiling process at high temperature (Sari *et al*, 2014). The aim of the experimentations was to evaluate the metal mobilization induced by citric acid from pineapple extract when samples were soaking and boiled until 45 minute.

**Article History:**

Received 24 January 2017; revised 27 January 2017; accepted 25 November 2017

<http://doi.org/10.26554/sti.2017.2.3.59-63>




Figure 1. Reduction of lead (Pb) in Juaro fish




Figure 2. Reduction of Hg in Juaro fish

## EXPERIMENTAL SECTION

### Experimental site

The experiment was performed in food and beverage analysis laboratory, faculty of health science, musi charitas catholic university, Palembang, Indonesia. The instrument was used a water-bath, hotplate magnetic stirrer and glassware instrument.

### Sample preparations

Juaro fish was purchased from traditional market in Palembang, South Sumatera, Indonesia. Samples of juaro fish was washed and separated their shells and meat. Meat of juaro fish was soaked in a extract of citric acid at 50 %, 75 %, 100 % for 15 min, 30 min, 45 min at 25 °C, 50 °C, 100 °C.

### Heavy metals analysis

The heavy metals such as Pb and Hg at juaro fish meat were analyzed by atomic absorption spectrophotometer (AAS) shimadzu AA 7000 in industrial research and standardization laboratory Palembang and absorption spectrophotometer (AAS) shimadzu AA 2800 in environmental health and disease control laboratory Palembang. Sample (5 g) were dissolved in 25 ml  $H_2SO_{4(p)}$ , 20 ml  $HNO_{3(p)}$ , 20 ml  $HNO_3-HClO_4(1:1)$  then analyzed using AAS. For mercury analysis, Mercury Vaporizer Unit (HVU) was added to AAS.

### Statistical analysis

The normality value of multiple groups were analyzed by one-way analysis of variance (ANOVA) with  $p < 0.05$  was classified as statistically significant.

## RESULT AND DISCUSSION

### Pb Content in Juaro fish (*Pangasius polyuranodon*)

Heavy metals contamination has risen and become one of the most dangerous pollutants of environment pollution. Continuously lead metal is released into the environment both naturally and human activities. The bioaccumulation of Pb in food should be reduced by using chelating agent. Citric acid as a chelating agent with vary concentration from pineapple extract, soaking time and boiled temperature was used to reduce the level Pb in juaro fish.

The level of Pb studied were successfully reduced by citric acid and are presented in fig 1. The obtained results from AAS showed that Pb content in juaro fish before soaking and boiled treatment was  $0.02 \text{ mg.Kg}^{-1}$ . The value shows that the Pb levels in juaro fish under the permissible limit. Based on Indonesia National standard the permissible limits of Pb content was  $1 \text{ mg.Kg}^{-1}$ .

Fish are considered as an important food source of macro-nutrients (carbohydrates, protein, fatty acids, vitamin and poly-unsaturated fatty acids) and micro nutrients (copper, zinc, iron, selenium) for human health (Arulkumar *et al*, 2017). Bioaccumulation of lead and high toxicity even at very low dosage can cause serious dangers to human health (Tajarudin *et al*, 2016). Lead (Pb) in juaro fish successfully remove above 80 % after soaked and boiled with 100 % pineapple extract for 45 min at 100 °C.

Citric acid was effective for the removal lead ions which is likely due to its  $C=C$  bond (Jiang *et al*, 2017). Removal effect of toxic metals among organic acids more effective at acidic condition. (Pitsari *et al*, 2016) Citric acid has large pka ( $pka = 3,15$ ) provide anions to complex with metals (doors, 2005). The statistical analysis showed that each treatment was significantly different and suggest that the higher boiled temperature and longer period of treatment time at any concentration of citric acid can increase the removal of Pb out from Juaro fish.

### Hg content in juaro fish (*Pangasius polyuranodon*)

Citric acid with varying concentration of pineapple extract, soaking time and boiling temperature was used to level of Hg in juaro fish. The levels of Hg studied were successfully reduced by Citric acid presented in fig 2. The Hg level in juaro fish from AAS instrument before soaking dan boiling treatment was  $0.011 \text{ mg.Kg}^{-1}$ . The value shows that the Hg levels in juaro fish under the permissible limit. Although below the permissible limit, contamination of Hg must be concern due to bioaccumulation, neurotoxicity and carcinogenicity (Jomova, 2010).

Heavy metals especially mercury has contaminated aquatic system (Malvandi, 2017), seafood species (Morgano *et al*, 2013), croaker fish (Musa *et al*, 2016), horseshoe crab (Bakker *et al*, 2017) and fish (Arulkumar *et al*, 2017). The mercury accumulated in body upon the consumption of contaminated fish. The adverse effect of Hg are associated with development abnormalities children (Parashor, 2013). Mercury in juaro fish successfully removed above 80 % after soaked and boiled with 100 % pineapple extract for 45 min at 100 °C.

The statistically analysis showed that each treatment was sig-

nificantly different and suggest that higher boiled temperature and longer period of treatment at any concentration of citric acid can increase removal of Hg out from juaro fish. It was indicates that an higher temperature related to increase the offered energy to separate the chemical bond of mercury metals in the materials and longer period of treatment time will give enough time for chelating agent to chelate with the heavy metals and extracted out from the juaro fish (Azele *et al*, 2014; Gzar *et al*, 2014)

## CONCLUSION

Citric acid was able to chelate the lead and mercury metals in juaro fish. The increasing of boiled temperature and longer period of treatment at any concentration of citric acid from pineapple extract can increase of removal heavy metals (Pb and Hg) in juaro fish. The highest decreasing of Pb and H level up to 80 % when samples were boiled in pineapple extract for 45 min at 100 °C. Based on statistical data, the sig value of Pb and Hg 0.000 and 0.04, respectively. Which showed the variation of pineapple extract have an effect on the decrease of Pb and Hg.

## ACKNOWLEDGEMENT

The authors are very thankful and appreciates to Indonesian Directorate General of Higher Education (DIKTI) for the financial support granted based on research of beginner lecturer.

## REFERENCES

- Aderholt, M; Vogilien, D.L; Kother, M., greipson, S. 2017. Phytoextraction of contaminated urban soils by *panicum virgatum* L enhanced with applications of a plant growth regulator (BAP) and citric acid. *Chemosphere* 175(2017) 85-96.
- Alloway, B.J., 1995. Heavy metals in soils. *Balckie Academic and professional*. 4-15
- Ahmed, M.K., Shaheen, N., Islam, M.S., Habibullah- Al-mamum M., Islam, S., Mohiduzaman, M., Bhattacharjee, L., 2015. Dietary intake of trace elements from highly consumed cultured fish (*labeo rohita*, *pangasius* and *oreochromis mossambicus*) and human health risk implications in Bangladesh. *Chemosphere* 128, 284-292.
- Arulkumar, A., Paramasivan, S., Rajaram., R. 2017. Toxic heavy metals in commercially important food fishes collected from palkbay southeast india. *Marine pollution bulletin* 3. <http://dx.doi.org/10.106/j.marpolbul.2017.03.045>
- Azelee, I.W., Ismail, R., Ali, R., Bakat, W.A. 2014. Chelation technique for the removals of heavy metals (As, Pb, Cd and Ni) from green mussel (*perna veridis*). *Indian Jurnal of geo-marine sciences* 43(3). 372-376
- Bakker, A.K., Dutton, J., Sclafani, M., Santangelo, N. 2017. Accumulation of nonessential trace elements (Ag, As, Cd, Cr, Hg and Pb) in atlantic horseshoe crab (*limulus polyphemus*) early life stages. *Science of total environment* 596-597 (2017) 69-78.
- Doors, S., 2005. Organic acids. In: Davidson, P.M., Sofos, J.N., Branen, A.L. (Eds.), *Antimicrobials in Food*. CRC Press, Boca Raton, FL, 91-142 (Taylor & Francis Group).
- Eddy S., Setiawan AA. Emilia I, Suheryanto. 2012. Bioaccumulation of mercury in various eco compartment of musi river Palembang. *Pekerti report PGRI-Sriwijaya University*
- Gzar, H., Abdul-hameed, A.S., Yahya, A.Y., 2014. Extraction of lead, cadmium and nikel from contaminated soil using acetic acid. *Open jurnal of soil science* 4. 207-214
- Goldstein, G.W. 1992. Neurological concepts of lead poisoning in children. *Pediatr Ann* 21 (6). 384-388.
- Hikmawati, A dan Lilis, S. 2006. Mobilization of mercury levels in Tuna Fish with soaking treatment of lime solution. Vol 3. Hal 67-76
- Jiang, J., Yang, M., Gao., Wang, J., Li, D., Li, T., 2017. Removal of toxic metals from vanadium-contaminated soils using a washing method : Reagent selection and parameter optimization. *Journal Chemosphere*. <http://dx.doi.org/10.1016/j.chemosphere.2017.03.116>
- Jomova, K., Valko, M. 2010. Advances in metal-induced oxidative stress and human disease. *Toxicology* 283. 65-67
- Kim, K.H., Kabir, E., Jahan, S.A., 2016. A review on the distribution of high in the environment and its human healths impact. *Jurnal of hazardous material*. Volume 306. pp 376-385. <https://doi.org/10.1016/j.jhazmat.2015.11.031>
- Liu, J., Yin, P., Chen, B., Gao, F., Song, H., Li, M., 2016. Distribution and contaminations assessment oh heavy metals in surface sediment of luanhe river estuary, north-west of bohai sea. *Pollution Bulletin*. <http://dx.doi.org/10.1016/j.marpolbul.2016.05.020>.
- Malvandi., H. 2017. Preliminary evaluation of heavy metals contamination in zarrin-gol river sediment. *Iran. Marine pollution bulletin*. <http://dx.doi.org/10.1016/j.marpolbul.2017.02.035>
- Morgano, M.A., Rabonato, L.C., Milani, R.F., Miyagusku, L., Quintaes, K.D., 2014. As, Cd, Cr, Pb and Hg in seafood species used for sashimi and evaluation of dietary exposure. *Jurnal food control*. 36 (2014). 24-29.
- Musa, A.M., Oyediji., D.E., Bello., M.S. 2016. Bioaccumulation of heavy metals in some tissues of croaker fish from oil spilled rivers of Niger delta region, Nigeria. *Asian Pacific journal of tropical biomedicine*.
- Parashor, P., Prasad, F.M. 2013. Study of heavy metals accumulation in sewage irrigated vegetables in different regions of agra district India. *Open.J.Soil.Sci.* 3 1-8.
- Pitsari, S., Tsoufakis, E., Loizidou, M. 2016. Enhanced lead adsorption by unbleached newspaper pulp modified with citric acid. *Environmental science and Technology Athens University. Chemical Engineering Journal*.
- Sanders, T., Liu, Y., Buchner, V., Tochounwou, P.B. 2009. Neurotoxic effects and biomarkers of lead exposure : a review. *Review Environment Health* 24. 15-45.
- Sari, K.A., Riyadi, P.H., Anggo.A.D. 2014. The effect of boiling temperature and lime (*Citrus aurantifolia*) solution on lead and cadmium levels on blood clams (*Anadara granosa*). *Oceanografi faculty of Diponegoro University. Biotechnology journal* vol 3 number 2. 1-10
- Shakour, M.B., All, S., Hameed, A., Farid, M., Hussain, S., Yasmeen, T., Najeeb, V., Bharwara, S.A., Abbasi, G.H., 2014. Citric acid improves lead (Pb) phtoextraction in *brassica napus* / by mitigating Pb-induced morphological and biochemical damages. *Ecotoxicology and environment safety* 109 (2014) 38-47. <http://dx.doi.org/10.1016/j.ecoenv.2014.07.033>
- Tadjuradi., Zadeh, A., Fendowski, S.M., Zare, R., Ahmad, B. 2016. Highly efficient ultrasonic assisted removal of Hg(II) ions on grapheme oxide modified with 2-pyridinecarboxaldehyde thiosemi carbazone : Adsorption isotherms and kinetics studies. *Ultrasonics sonochemistry*. Iran University of science and technology.

- Turaga, R.M.R., Howarth, R.B., Borsuk, M.E. Perceptions of mercury risk and its management human and ecological risk assessment. An International Journal.
- Ulfah, S., Fida R., Raharjo. 2014. Mobilization of lead content in *Mystus nigriceps* at Kali rivers of Surabaya using extract of pineapple skin. LenteraBio. 2(1) : 103-108